



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE CONDITION THAT A LINEAR TOTAL DIFFERENTIAL EQUATION BE INTEGRABLE.

BY E. O. LOVETT.

IN order that the linear partial differential equations

$$(1) \quad \Omega_1 f \equiv \xi_1(x, y, z) \frac{\partial f}{\partial x} + \eta_1(x, y, z) \frac{\partial f}{\partial y} + \zeta_1(x, y, z) \frac{\partial f}{\partial z} = 0,$$

$$(2) \quad \Omega_2 f \equiv \xi_2(x, y, z) \frac{\partial f}{\partial x} + \eta_2(x, y, z) \frac{\partial f}{\partial y} + \zeta_2(x, y, z) \frac{\partial f}{\partial z} = 0,$$

have a common solution it is both necessary and sufficient that a relation of the form

$$(\Omega_1, \Omega_2) \equiv \Omega_1(\Omega_2 f) - \Omega_2(\Omega_1 f) = \lambda_1(x, y, z) \Omega_1 f + \lambda_2(x, y, z) \Omega_2 f$$

exist identically, whatever the function $f(x, y, z)$ may be.*

If the above equations have the forms

$$(3) \quad X_1 f \equiv \frac{\partial f}{\partial x} + \theta_1(x, y, z) \frac{\partial f}{\partial z} = 0,$$

$$(4) \quad X_2 f \equiv \frac{\partial f}{\partial y} + \theta_2(x, y, z) \frac{\partial f}{\partial z} = 0,$$

the expression (X_1, X_2) will have no terms in $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$, and λ_1, λ_2 accordingly become zero, that is

$$(X_1, X_2) \equiv X_1(X_2 f) - X_2(X_1 f) = 0,$$

is the necessary and sufficient condition that (3) and (4) have a common solution.

In order that the linear total differential equation

$$(5) \quad P(x, y, z) dx + Q(x, y, z) dy + R(x, y, z) dz = 0$$

* See Lie-Scheffers, *Differentialgleichungen mit bekannten infinitesimalen Transformationen*, chap. 10, § 3; Leipzig, 1891.

have an integral it is both necessary and sufficient that the linear partial differential equations

$$Uf \equiv \frac{\partial f}{\partial x} - \frac{P}{R} \frac{\partial f}{\partial z} = 0, \quad Vf \equiv \frac{\partial f}{\partial y} - \frac{Q}{R} \frac{\partial f}{\partial z} = 0$$

have a common solution.

Hence in order that the linear total differential equation (5) be integrable it is both necessary and sufficient that

$$(U, V) = 0;$$

or, in developed form

$$\frac{1}{R^2} \left\{ P(Q_z - R_y) + Q(R_x - P_z) + R(P_y - Q_x) \right\} \frac{\partial f}{\partial z} = 0;$$

that is

$$P(Q_z - R_y) + Q(R_x - P_z) + R(P_y - Q_x) = 0.$$

The above method admits of immediate extension to the case of more than three independent variables

$$\sum_1^n P_i(x_1, x_2, \dots, x_n) dx_i = 0.$$

PRINCETON, NEW JERSEY, APRIL, 1900.